

# Bering Sea Project

**Bering Ecosystem Study (BEST) +  
Bering Sea Integrated Ecosystem Research Program (BSIERP)**



North Pacific Research Board  
Board of Directors meeting  
Anchorage, Alaska  
April 30, 2014

Mike Sigler (NOAA)  
(on behalf of the larger program)



# Program scope and chronology




- **2007 – 2010** Field Work
- **2011 – 2013** Synthesis
- **24,205** person-days of fieldwork
- **136** publications to date

# Outline

- Prologue: Seasonal ice and the cold pool
- Chapter 1: Why did pollock abundance decline then rebound in the last decade?
- Chapter 2: Location matters for fur seals and fishermen
- Chapter 3: The eastern Bering Sea in the future
- The benefits of an integrated program



# Prologue: Icy winters occur when winds are from the north and Arctic in origin





Satellite image climate.gov, wind image jacksmumontherun.wordpress.com



# Icy winters increase the size of the 'cold pool' ( $<2\text{ }^{\circ}\text{C}$ )

WARM YEAR

COLD YEAR




Lauth

# Chapter 1


## EXPLAIN THIS:

Walleye pollock abundance dramatically fell in the early 2000's, leading to a 40% drop in the quota for the largest single fishery in the US, and then rebounded.


Age-1 number (millions)



Age-3+  
Biomass  
(thousands t)



The amount of primary production available for copepods and krill is similar in warm and cold years



✱ 1979-1981 (Whitledge et al., 1986)

□ 1997 (Stockwell et al., 2001)

△ 1983 (Hansell et al., 1993)

○ 2008-2009 (BEST)

◇ 2004 (EcoFOCI / NOAA)


... and does not appear to limit production of copepod and krill, which are prey for age-0 pollock

Mordy, Cokelet, Ladd, Menzies, Proctor, Stabenow, Wisegarver

Campbell, Ashjian, Lessard, Liu, Zhai, Zeeman, Eisner, Gann, Mordy, Moran, Lomas, Gibson






# Bloom timing matches copepod egg production in cold years






# Bloom timing matches copepod egg production in cold years but not warm years

*Calanus* egg production


-  Spring open water blooms
-  Fall open water blooms
-  Ice assoc. blooms



# Ice algae likely enhances copepod reproduction



Gradinger, Bluhm, Iken,  
Weems




Higher ingestion rate when feeding on ice algae than water column phytoplankton

Campbell, Lessard, Ashjian,  
Durbin, Ryneason, Casas




Copepods and krill are more abundant in cold years: This contradicted our expectation (strike 1!)

Large zooplankton abundance




As a consequence, age-0 pollock consume richer diets in cold years, better preparing them for their first winter...



... and enhancing survivorship.

Survivorship Anomaly




Heintz et al.




## EXPLANATION:

Due to bloom timing, large crustacean zooplankton benefit from icy winters, providing prey for age-0 pollock to enter their first winter fat (and happy?)

### Age-1 number (millions)



Age-3+  
Biomass  
(thousands t)




# Chapter 2: Location matters for fur seals and fishermen




We predicted fishermen would travel farther north in warm years, but instead the opposite occurred (strike 2!)

Pollock catcher/  
processor fleet,  
center of fishing  
effort:

In summer, a  
northward and off-  
shelf shift correlated  
with colder  
conditions and larger  
cold pool




# Even with negligible temperature-related shift, some cost effects can occur




Summer/fall cod longline fishery: Vessels traveled farther (29 vs 20 km/ton catch) and set their gear more often during a trip (39 vs 33) in warm vs. cold years.



# Population trends differ among locations





# Population trends differ among locations



# Foraging locations - closer to home is better

- Trip lengths shorter for fur seals and murrelets at Bogoslof than Pribilofs
- Energy content of diet lower at Pribilofs than Bogoslof because of species consumed


Trites, Battaile, Benoit-Bird, Harding, Heppell, Irons, Kitaysky, Kuletz, Paredes, Renner, Roby



Kernel density use estimates for northern fur seals, blacklegged kittiwakes, and thickbilled murrelets tagged on St. Paul Island (purple), St. George Island (green) and Bogoslof Island (blue) in 2009. Isoleths are 50, 75, and 95 % use contours with darker colors indicating higher use areas.

Trends observed by communities (local and traditional knowledge, subsistence harvests):

- Location differences between the south (many species in decline) and the north (a productive ecosystem)
- Patterns are consistent with the northern Bering Sea remaining icy during winter and spring and the southeastern Bering Sea more affected by changes in sea ice extent




Huntington, Braem, Brown, Hunn, Krieg, Lestenkof, Noongwook, Sepez, Sigler, Wiese, Zavadil




# Chapter 3: The eastern Bering Sea in the future

# Future ocean conditions: The north will remain cold and dark

Warm years




Cold years



The anomalies of sea-ice coverage during March and April during warm years (2001-2005, left) and cold years (2007-2010, right) (Stabeno, Farley, Kachel, Moore, Mordy, Napp, Overland, Pinchuk, Sigler)


Subarctic fish will not expand into the northern Bering Sea shelf, which contradicts our expectation when the program started (strike 3!)



The average number of days in which sea-ice was present in March and April during 2001-2010.

Stabeno, Farley, Kachel, Moore, Mordy, Napp, Overland, Pinchuk, Sigler, Hollowed, Barbeaux, Cokelet, Kotwicki, Ressler, Spital, Wilson

# Models also forecast that the north will remain cold and dark



Zhang and Banas







# Forecast fish abundance, climate effects differ



### Temperature







### Inshore transport



Mueter, Bond, Ianelli, Hollowed

Wilderbuer, Stockhausen, Bond

# Forecast fish abundance, climate effects differ



Mueter, Bond, Ianelli, Hollowed

Wilderbuer, Stockhausen, Bond

# The benefits of an integrated ecosystem research program


- Management implications
  - understanding “why” helps stakeholders
- Deep and broad publication set
- Formation of new teams and collaborations
- Results achieved by this integrated program that likely would not have been accomplished by a series of individual projects (3 examples)



Due to bloom timing, large crustacean zooplankton benefit from icy winters, providing prey for age-0 pollock to enter their first winter fat





ICE



COLD YEAR






Age-1 number (millions)



Low energy demand





# Prey closer to colonies and more energy dense at Bogoslof compared to Pribilof islands



Trites, Battaile, Benoit-Bird, Friday, Harding, Heppell, Hoover, Irons, Jones, Kitaysky, Kuletz, McIntosh, Mueter, Nordstrom, Orben, Paredes, Renner, Ressler, Roby, Sigler, Suryan, Waluk, Wilson, Young, Zerbini



# The northern Bering Sea will stay cold for the foreseeable future and subarctic fish will not expand into there



Banas, Barbeaux, Bond, Cokelet, Curchitser, Farley, Gibson, Hedstrom, Hermann, Hollowed, Kachel, Kotwicki, Moore, Mordy, Napp, Overland, Pinchuk, Ressler, Sigler, Spital, Stabeno, Wilson, Zhang

