Marine Trade Wind Cumulus


Status of CVI Measurements from RICO

RICO Analysis Meeting 28 June 2005


Scientific Questions

- What are the nuclei of droplets in maritime cumulus? Is there evidence of pollution in this remote area?
- What are the nuclei of the largest droplets, which initiate precipitation? What are the giant nuclei concentrations and sources?
- What are the relationships between particle size, composition and critical supersaturation, and do they agree with theory? (organics)
- How do these compare with polluted mixed-phase clouds (AIRS-II) and California stratus (DYCOMS-II)?

Counterflow Virtual Impactor (CVI)




- Drops and ice crystals are impacted into dry nitrogen gas and evaporated.
 Interstitial particles and gases excluded.
- Measure volatile gases (water vapor) and non-volatile residual particles
- Change min. size drop sampled to investigate drop size dependence
- Also used as "giant nuclei" sampler outside of cloud




CVI Measurements


- CVI operated well on all flights; major problem is breakup of droplets
 50-100 microns when sampling--may produce > 1 residual nucleus
- 2 criteria for breakup screening of electron microscope samples:
 [(CVI N)/(FSSP N)] > 1.25 and 2DP LWC > 0.0000




- 84 samples screened, 19 (23%) definite pass, 17 (20%) possible pass
- Only about 2/3 flights screened (REU student)
- Each in-cloud sample has a filter and 4 impactor grids, corresponding below cloud total and giant nuclei sample, also some above cloud
- Plenty of samples to analyze! Starting now...

Data Example Flight with Little Precip









Northern Hem. marine environment is polluted, even away from major sources and in the tropics


Caribbean is Not Pristine






Interesting Things Noted: Aerosol Indirect Effect in Caribbean?


Below-cloud circles for R6 (16 Dec)




Strong effect on drop N



DYCOMS-II Indirect Effect






Twohy et al. 2005

Is it important for RICO? Need to stratify by updraft V

Interesting Things Noted New Particle Production in Outflow Region



Data Analysis Goals

(collaboration with ASU, DRI, NCAR)

- Analyze all high priority SEM and TEM samples
 quantitatively (below cloud, in cloud at low and high cut
 sizes if available), above and in cloud outflow regions)
- · Archive number and liquid water data for high priority cases
- Analyze Giant Nuclei samples from various samplers (CVI, TAS, UGN Stick) for concentration and composition
- Examine composition in greater detail via Electron Energy Loss Spectroscopy (organics)
- Compare ambient and cloud droplet residual CCN spectra and size distributions
- Return to Scientific Questions

Scientific Questions

- What are the nuclei of droplets in maritime cumulus? Is there evidence of pollution in this remote area?
- What are the nuclei of the largest droplets, which initiate precipitation? What are the giant nuclei concentrations and sources?
- What are the relationships between particle size, composition and critical supersaturation, and do they agree with theory? (organics)
- How do these compare with polluted mixed-phase clouds (AIRS-II) and California stratus (DYCOMS-II)?